

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssptamym1652

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 SEP 09 ACD predicted properties enhanced in REGISTRY/ZREGISTRY  
NEWS 4 OCT 03 MATHDI removed from STN  
NEWS 5 OCT 04 CA/CAPplus-Canadian Intellectual Property Office (CIPO) added  
to core patent offices  
NEWS 6 OCT 13 New CAS Information Use Policies Effective October 17, 2005  
NEWS 7 OCT 17 STN(R) AnaVist(TM), Version 1.01, allows the export/download  
of CAPplus documents for use in third-party analysis and  
visualization tools  
NEWS 8 OCT 27 Free KWIC format extended in full-text databases  
NEWS 9 OCT 27 DIOGENES content streamlined  
NEWS 10 OCT 27 EPFULL enhanced with additional content  
NEWS 11 NOV 14 CA/CAPplus - Expanded coverage of German academic research  
NEWS 12 NOV 30 REGISTRY/ZREGISTRY on STN(R) enhanced with experimental  
spectral property data  
NEWS 13 DEC 05 CASREACT(R) - Over 10 million reactions available  
  
NEWS EXPRESS DECEMBER 02 CURRENT VERSION FOR WINDOWS IS V8.01,  
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 02 DECEMBER 2005.  
V8.0 USERS CAN OBTAIN THE UPGRADE TO V8.01 AT  
<http://download.cas.org/express/v8.0-Discover/>  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:21:08 ON 05 DEC 2005

=> file .mymstn

COST IN U.S. DOLLARS

SINCE FILE  
ENTRY

TOTAL  
SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'MEDLINE' ENTERED AT 12:21:42 ON 05 DEC 2005

FILE 'AGRICOLA' ENTERED AT 12:21:42 ON 05 DEC 2005

FILE 'JICST-EPLUS' ENTERED AT 12:21:42 ON 05 DEC 2005  
COPYRIGHT (C) 2005 Japan Science and Technology Agency (JST)

FILE 'BIOSIS' ENTERED AT 12:21:42 ON 05 DEC 2005  
Copyright (c) 2005 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 12:21:42 ON 05 DEC 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'LIFESCI' ENTERED AT 12:21:42 ON 05 DEC 2005  
COPYRIGHT (C) 2005 Cambridge Scientific Abstracts (CSA)

FILE 'BIOTECHNO' ENTERED AT 12:21:42 ON 05 DEC 2005  
COPYRIGHT (C) 2005 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'EMBASE' ENTERED AT 12:21:42 ON 05 DEC 2005  
Copyright (c) 2005 Elsevier B.V. All rights reserved.

=> s elongase  
L1 1349 ELONGASE

=> s elongate? (3w) fatty (w) acid  
L2 229 ELONGATE? (3W) FATTY (W) ACID

=>

=> s Oomycete  
L3 5863 OOMYCETE

=> s l1 and l2 and l3  
L4 0 L1 AND L2 AND L3

=> s l1 and l2  
L5 17 L1 AND L2

=> d ibib abs l5 1-17

L5 ANSWER 1 OF 17 MEDLINE on STN  
ACCESSION NUMBER: 2005551393 IN-PROCESS  
DOCUMENT NUMBER: PubMed ID: 15129327  
TITLE: Zebrafish cDNA encoding multifunctional Fatty Acid  
elongase involved in production of eicosapentaenoic  
(20:5n-3) and docosahexaenoic (22:6n-3) acids.  
AUTHOR: Agaba Morris; Tocher Douglas R; Dickson Cathryn A; Dick  
James R; Teale Alan J  
CORPORATE SOURCE: Institute of Aquaculture, University of Stirling, Stirling  
FK9 4LA, Scotland, UK.  
SOURCE: Marine biotechnology (New York, N.Y.), (2004 May-Jun) 6 (3)  
251-61. Electronic Publication: 2004-05-06.  
Journal code: 100892712. ISSN: 1436-2228.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: NONMEDLINE; IN-PROCESS; NONINDEXED; Priority Journals  
ENTRY DATE: Entered STN: 20051018  
Last Updated on STN: 20051020

AB Enzymes that increase the chain length of fatty acids are essential for

biosynthesis of highly unsaturated fatty acids. The gLELO gene encodes a protein involved in the elongation of polyunsaturated fatty acids in the fungus *Mortierella alpina*. A search of the GenBank database identified several expressed sequence tag sequences, including one obtained from zebrafish (*Danio rerio*), with high similarity to gLELO. The full-length transcript ZfELO, encoding a polypeptide of 291 amino acid residues, was isolated from zebrafish liver cDNA. The predicted amino acid sequence of the open reading frame shared high similarity with the **elongases** of *Caenorhabditis elegans* and human. When expressed in *Saccharomyces cerevisiae*, the zebrafish open reading frame conferred the ability to lengthen the chain of a range of C18, C20, and C22 polyunsaturated fatty acids, indicating not only that biosynthesis of 22:6n-3 from 18:3n-3 via a 24-carbon intermediate is feasible, but also that one **elongase** enzyme can perform all three elongation steps required. The zebrafish enzyme was also able to **elongate** monounsaturated and saturated **fatty acids**, and thus demonstrates a greater level of promiscuity in terms of substrate use than any **elongase** enzyme described previously.

L5 ANSWER 2 OF 17 MEDLINE on STN  
 ACCESSION NUMBER: 2005027689 MEDLINE  
 DOCUMENT NUMBER: PubMed ID: 15652569  
 TITLE: Biochemical characterization of **elongase** activity in corn (*Zea mays* L.) roots.  
 AUTHOR: Schreiber Lukas; Franke Rochus; Lessire Rene  
 CORPORATE SOURCE: Department of Ecophysiology, Institute of Cellular and Molecular Botany (IZMB), University of Bonn, Kirschallee 1, D-53115 Bonn, Germany.. lukas.schreiber@uni-bonn.de  
 SOURCE: Phytochemistry, (2005 Jan) 66 (2) 131-8.  
 Journal code: 0151434. ISSN: 0031-9422.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200504  
 ENTRY DATE: Entered STN: 20050119  
 Last Updated on STN: 20050412  
 Entered Medline: 20050411

AB Chemical analysis of 4-day-old corn (*Zea mays* L.) root cell walls revealed that the lipophilic biopolymer suberin forms an important constituent of rhizodermal and hypodermal cell walls. Identified aliphatic monomers had chain lengths ranging from C16 to C26 and they belonged to 5 substance classes (omega-hydroxycarboxylic acids, 1,omega-dicarboxylic acids, 2-hydroxycarboxylic acids, carboxylic acids and alcohols) by which suberin is characterized. Biochemical experiments proved the occurrence of **elongase** activities in corn roots. Highest enzymatic activities were found in corn root microsomes, and major products synthesized by root **elongases** were **elongated fatty acids** with chain lengths ranging from C20 to C24. Preferred substrates of root **elongases** were acyl-CoAs of the chain length C18 and C20, whereas monounsaturated acyl-CoAs (C16:1 and C18:1) and acyl-CoAs of lower (C12-C16) and higher chain lengths (C22-C24) were rarely elongated. **Elongase** activities significantly decreased over the length (40 cm) of 10-day-old corn roots going from the young tip to the older base of the root. Thus, results presented here show the presence and activity of **elongases** in roots of plants.

L5 ANSWER 3 OF 17 MEDLINE on STN  
 ACCESSION NUMBER: 2004499304 MEDLINE  
 DOCUMENT NUMBER: PubMed ID: 15377762  
 TITLE: Biosynthesis of very-long-chain polyunsaturated fatty acids in transgenic oilseeds: constraints on their accumulation.  
 AUTHOR: Abbadi Amine; Domergue Frederic; Bauer Jorg; Napier Johnathan A; Welte Ruth; Zahringer Ulrich; Cirpus Petra;

Heinz Ernst  
 CORPORATE SOURCE: Institut fur Allgemeine Botanik, Universitat Hamburg, 22609  
 Hamburg, Germany. abbadi@botanik.uni-hamburg.de.  
 <abbadi@botanik.uni-hamburg.de>  
 SOURCE: Plant cell, (2004 Oct) 16 (10) 2734-48. Electronic  
 Publication: 2004-09-17.  
 Journal code: 9208688. ISSN: 1040-4651.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 OTHER SOURCE: GENBANK-AF054824; GENBANK-AF428243; GENBANK-AJ222980;  
 GENBANK-AX035537; GENBANK-AY082392; GENBANK-AY082393;  
 GENBANK-U79010  
 ENTRY MONTH: 200503  
 ENTRY DATE: Entered STN: 20041007  
 Last Updated on STN: 20050323  
 Entered Medline: 20050322

AB Omega6- and omega3-polyunsaturated C20 fatty acids represent important  
 components of the human diet. A more regular consumption and an  
 accordingly sustainable source of these compounds are highly desirable.  
 In contrast with the very high levels to which industrial fatty acids have  
 to be enriched in plant oils for competitive use as chemical feedstocks,  
 much lower percentages of very-long-chain polyunsaturated fatty acids  
 (VLCPUFA) in edible plant oils would satisfy nutritional requirements.  
 Seed-specific expression in transgenic tobacco (*Nicotiana tabacum*) and  
 linseed (*Linum usitatissimum*) of cDNAs encoding fatty acyl-desaturases and  
 elongases, absent from all agronomically important plants,  
 resulted in the very high accumulation of Delta6-desaturated C18 fatty  
 acids and up to 5% of C20 polyunsaturated fatty acids, including  
 arachidonic and eicosapentaenoic acid. Detailed lipid analyses of  
 developing seeds from transgenic plants were interpreted as indicating  
 that, after desaturation on phosphatidylcholine, Delta6-desaturated  
 products are immediately channeled to the triacylglycerols and effectively  
 bypass the acyl-CoA pool. Thus, the lack of available Delta6-desaturated  
 acyl-CoA substrates in the acyl-CoA pool limits the synthesis of  
 elongated C20 fatty acids and disrupts the  
 alternating sequence of lipid-linked desaturations and acyl-CoA dependent  
 elongations. As well as the successful production of VLCPUFA in  
 transgenic oilseeds and the identification of constraints on their  
 accumulation, our results indicate alternative strategies to circumvent  
 this bottleneck.

L5 ANSWER 4 OF 17 MEDLINE on STN  
 ACCESSION NUMBER: 2001695342 MEDLINE  
 DOCUMENT NUMBER: PubMed ID: 11743108  
 TITLE: A condensing enzyme from the seeds of *Lesquerella fendleri*  
 that specifically elongates hydroxy fatty  
 acids.  
 AUTHOR: Moon H; Smith M A; Kunst L  
 CORPORATE SOURCE: Department of Botany, University of British Columbia,  
 Vancouver, British Columbia V6T 1Z4, Canada.  
 SOURCE: Plant physiology, (2001 Dec) 127 (4) 1635-43.  
 Journal code: 0401224. ISSN: 0032-0889.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200204  
 ENTRY DATE: Entered STN: 20011217  
 Last Updated on STN: 20020417  
 Entered Medline: 20020416

AB *Lesquerella fendleri* seed oil contains up to 60% hydroxy fatty acids,  
 nearly all of which is the 20-carbon hydroxy fatty acid lesquerolic acid

(D-14-hydroxyeicos-cis-11-enoic acid). Previous work suggested that lesquerolic acid in *L. fendleri* was formed by the elongation of the 18-carbon hydroxy fatty acid, ricinoleic acid. To identify a gene encoding the enzyme involved in hydroxy fatty acid elongation, an *L. fendleri* genomic DNA library was screened using the coding region of the Arabidopsis Fatty Acid Elongation1 gene as a probe. A gene, LfKCS3, with a high sequence similarity to known very long-chain fatty acid condensing enzymes, was isolated. LfKCS3 has a 2,062-bp open reading frame interrupted by two introns, which encodes a polypeptide of 496 amino acids. LfKCS3 transcripts accumulated only in the embryos of *L. fendleri* and first appeared in the early stages of development. Fusion of the LfKCS3 promoter to the uidA reporter gene and expression in transgenic Arabidopsis resulted in a high level of beta-glucuronidase activity exclusively in developing embryos. Seeds of Arabidopsis plants transformed with LfKCS3 showed no change in their very long-chain fatty acid content. However, when these Arabidopsis plants were crossed with the transgenic plants expressing the castor oleate 12-hydroxylase, significant amounts of 20-carbon hydroxy fatty acids accumulated in the seed, indicating that the LfKCS3 condensing enzyme specifically catalyzes elongation of 18-carbon hydroxy fatty acids.

- L5 ANSWER 5 OF 17 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.  
(2005) on STN

ACCESSION NUMBER: 2005:15681 AGRICOLA  
DOCUMENT NUMBER: IND43672333  
TITLE: Biochemical characterization of elongase activity in corn (*Zea mays* L.) roots.  
AUTHOR(S): Schreiber, L.; Franke, R.; Lessire, R.  
AVAILABILITY: DNAL (450 P5622)  
SOURCE: Phytochemistry, 2005 Jan. Vol. 66, no. 2 p. 131-138  
ISSN: 0031-9422  
NOTE: Includes references  
DOCUMENT TYPE: Article  
FILE SEGMENT: Non-US  
LANGUAGE: English

AB Chemical analysis of 4-day-old corn (*Zea mays* L.) root cell walls revealed that the lipophilic biopolymer suberin forms an important constituent of rhizodermal and hypodermal cell walls. Identified aliphatic monomers had chain lengths ranging from C16 to C26 and they belonged to 5 substance classes (omega-hydroxycarboxylic acids, 1,omega-dicarboxylic acids, 2-hydroxycarboxylic acids, carboxylic acids and alcohols) by which suberin is characterized. Biochemical experiments proved the occurrence of elongase activities in corn roots. Highest enzymatic activities were found in corn root microsomes, and major products synthesized by root elongases were elongated fatty acids with chain lengths ranging from C20 to C24. Preferred substrates of root elongases were acyl-CoAs of the chain length C18 and C20, whereas monounsaturated acyl-CoAs (C16:1 and C18:1) and acyl-CoAs of lower (C12-C16) and higher chain lengths (C22-C24) were rarely elongated. Elongase activities significantly decreased over the length (40 cm) of 10-day-old corn roots going from the young tip to the older base of the root. Thus, results presented here show the presence and activity of elongases in roots of plants.

- L5 ANSWER 6 OF 17 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.  
(2005) on STN

ACCESSION NUMBER: 1998:42513 AGRICOLA  
DOCUMENT NUMBER: IND21233827  
TITLE: Fatty acid and wax biosynthesis in susceptible and triallate-resistant *Avena fatua* L.

AUTHOR(S): Kern, A.J.; Jackson, L.L.; Dyer, W.E.  
AVAILABILITY: DNAL (SB951.P47)  
SOURCE: Pesticide science, Sept 1997. Vol. 51, No. 1. p. 21-26  
Publisher: Chichester, West Sussex : John Wiley and  
Sons Limited.  
CODEN: PSSCBG; ISSN: 0031-613X  
NOTE: Includes references  
PUB. COUNTRY: England; United Kingdom  
DOCUMENT TYPE: Article  
FILE SEGMENT: Non-U.S. Imprint other than FAO  
LANGUAGE: English

AB The recent characterization of triallate-resistant lines of wild oat (*Avena fatua* L.) deficient in triallate sulfoxidation provides an experimental system to investigate and differentiate the effects of triallate and triallate sulfoxide on wax and lipid biosynthesis. Greenhouse applications of triallate dramatically reduced epicuticular wax deposition in susceptible (S) but not resistant (R) wild oats. Triallate treatment had no effect on in-vivo concentrations of C12 to C26 fatty acids and fatty alcohols in R plants, while **elongated fatty acid** fractions (C > 18) were significantly reduced in S plants. In contrast, treatment with triallate sulfoxide reduced in-vivo concentrations of **elongated fatty acids** equally in R and S, supporting the hypothesis that triallate sulfoxide is more inhibitory than triallate towards fatty acid **elongases**. Although de-novo synthesis of short-chain fatty acids was not affected by triallate or triallate sulfoxide in R or S plants, synthesis of **elongated fatty acid** fractions was dramatically reduced in S plants by triallate. Fatty acid biosynthesis in R and S plants was equally sensitive to triallate sulfoxide. The results support the idea that in-vivo triallate sulfoxidation is necessary for herbicidal activity, and confirm that reduced rates of triallate sulfoxidation confer resistance in R wild oats.

L5 ANSWER 7 OF 17 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN  
ACCESSION NUMBER: 2005:176735 BIOSIS  
DOCUMENT NUMBER: PREV200500174426  
TITLE: Biochemical characterization of **elongase** activity  
in corn (*Zea mays* L.) roots.  
AUTHOR(S): Schreiber, Lukas [Reprint Author]; Franke, Rochus; Lessire, Ren  
CORPORATE SOURCE: Dept EcophysiolInst Mol and Cellular BiolIZMT, Univ Bonn, Kirschallee 1, D-53115, Bonn, Germany  
lukas.schreiber@uni-bonn.de  
SOURCE: Phytochemistry (Amsterdam), (January 2005) Vol. 66, No. 2, pp. 131-138. print.  
ISSN: 0031-9422 (ISSN print).  
DOCUMENT TYPE: Article  
LANGUAGE: English  
ENTRY DATE: Entered STN: 4 May 2005  
Last Updated on STN: 4 May 2005

AB Chemical analysis of 4-day-old corn (*Zea mays* L.) root cell walls revealed that the lipophilic biopolymer suberin forms an important constituent of rhizodermal and hypodermal cell walls. Identified aliphatic monomers had chain lengths ranging from C16 to C26 and they belonged to 5 substance classes ( $\omega$ -hydroxycarboxylic acids, 1, $\omega$ -dicarboxylic acids, 2-hydroxycarboxylic acids, carboxylic acids and alcohols) by which suberin is characterized. Biochemical experiments proved the occurrence of **elongase** activities in corn roots. Highest enzymatic activities were found in corn root microsomes, and major products synthesized by root **elongases** were **elongated fatty acids** with chain lengths ranging from C20 to C24. Preferred substrates of root **elongases** were acyl-CoAs of the chain length C18 and C20, whereas monounsaturated acyl-CoAs (C16:1 and C18:1) and acyl-CoAs of lower (C12-C16) and higher chain lengths (C22-C24) were rarely elongated.

Elongase activities significantly decreased over the length (40 cm) of 10-day-old corn roots going from the young tip to the older base of the root. Thus, results presented here show the presence and activity of elongases in roots of plants. Copyright 2004 Elsevier Ltd. All rights reserved.

L5 ANSWER 8 OF 17 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN  
ACCESSION NUMBER: 2005:42417 BIOSIS  
DOCUMENT NUMBER: PREV200500041515  
TITLE: Biosynthesis of very-long-chain polyunsaturated fatty acids in transgenic oilseeds: Constraints on their accumulation.  
AUTHOR(S): Abbadi, Amine [Reprint Author]; Domergue, Frederic; Bauer, Joerg; Napier, Johnathan A.; Welte, Ruth; Zaehring, Ulrich; Cirpus, Petra; Heinz, Ernst  
CORPORATE SOURCE: Inst Allgemeine Bot, Univ Hamburg, H-22609, Hamburg, Germany  
abbadi@botanik.uni-hamburg.de; eheinz@botanik.uni-hamburg.de  
SOURCE: Plant Cell, (October 2004) Vol. 16, No. 10, pp. 2734-2748. print.  
CODEN: PLCEEW. ISSN: 1040-4651.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
ENTRY DATE: Entered STN: 26 Jan 2005  
Last Updated on STN: 26 Jan 2005

AB omega6- and omega3-polyunsaturated C20 fatty acids represent important components of the human diet. A more regular consumption and an accordingly sustainable source of these compounds are highly desirable. In contrast with the very high levels to which industrial fatty acids have to be enriched in plant oils for competitive use as chemical feedstocks, much lower percentages of very-long-chain polyunsaturated fatty acids (VLCPUFA) in edible plant oils would satisfy nutritional requirements. Seed-specific expression in transgenic tobacco (*Nicotiana tabacum*) and linseed (*Linum usitatissimum*) of cDNAs encoding fatty acyl-desaturases and elongases, absent from all agronomically important plants, resulted in the very high accumulation of DELTA6-desaturated C18 fatty acids and up to 5% of C20 polyunsaturated fatty acids, including arachidonic and eicosapentaenoic acid. Detailed lipid analyses of developing seeds from transgenic plants were interpreted as indicating that, after desaturation on phosphatidylcholine, DELTA6-desaturated products are immediately channeled to the triacylglycerols and effectively bypass the acyl-CoA pool. Thus, the lack of available DELTA6-desaturated acyl-CoA substrates in the acyl-CoA pool limits the synthesis of elongated C20 fatty acids and disrupts the alternating sequence of lipid-linked desaturations and acyl-CoA dependent elongations. As well as the successful production of VLCPUFA in transgenic oilseeds and the identification of constraints on their accumulation, our results indicate alternative strategies to circumvent this bottleneck.

L5 ANSWER 9 OF 17 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN  
ACCESSION NUMBER: 2004:348186 BIOSIS  
DOCUMENT NUMBER: PREV200400349937  
TITLE: Zebrafish cDNA encoding multifunctional fatty acid elongase involved in production of eicosapentaenoic (20:5n-3) and docosahexaenoic (22:6n-3) acids.  
AUTHOR(S): Agaba, Morris; Tocher, Douglas R. [Reprint Author]; Dickson, Cathryn A.; Dick, James R.; Teale, Alan J.  
CORPORATE SOURCE: Inst Aquaculture, Univ Stirling, Stirling, FK9 4LA, Scotland  
d.r.tocher@stir.ac.uk  
SOURCE: Marine Biotechnology (New York), (May 2004) Vol. 6, No. 3, pp. 251-261. print.  
ISSN: 1436-2228 (ISSN print).

DOCUMENT TYPE: Article  
LANGUAGE: English  
ENTRY DATE: Entered STN: 18 Aug 2004  
Last Updated on STN: 18 Aug 2004

AB Enzymes that increase the chain length of fatty acids are essential for biosynthesis of highly unsaturated fatty acids. The gLELO gene encodes a protein involved in the elongation of polyunsaturated fatty acids in the fungus *Mortierella alpina*. A search of the GenBank database identified several expressed sequence tag sequences, including one obtained from zebrafish (*Danio rerio*), with high similarity to gLELO. The full-length transcript ZfELO, encoding a polypeptide of 291 amino acid residues, was isolated from zebrafish liver cDNA. The predicted amino acid sequence of the open reading frame shared high similarity with the **elongases** of *Caenorhabditis elegans* and human. When expressed in *Saccharomyces cerevisiae*, the zebrafish open reading frame conferred the ability to lengthen the chain of a range of C18, C20, and C22 polyunsaturated fatty acids, indicating not only that biosynthesis of 22:6n-3 from 18:3n-3 via a 24-carbon intermediate is feasible, but also that one **elongase** enzyme can perform all three elongation steps required. The zebrafish enzyme was also able to **elongate** monounsaturated and saturated **fatty acids**, and thus demonstrates a greater level of promiscuity in terms of substrate use than any **elongase** enzyme described previously.

L5 ANSWER 10 OF 17 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN

ACCESSION NUMBER: 1997:459396 BIOSIS  
DOCUMENT NUMBER: PREV199799758599  
TITLE: Fatty acid and wax biosynthesis in susceptible and triallate-resistant *Avena fatua* L.  
AUTHOR(S): Kern, Anthony J. [Reprint author]; Jackson, Larry L.; Dyer, William E. [Reprint author]  
CORPORATE SOURCE: Dep. Plant Soil Environ. Sci., Montana State Univ., Bozeman, MT 59717-0312, USA  
SOURCE: Pesticide Science, (1997) Vol. 51, No. 1, pp. 21-26.  
CODEN: PSSCBG. ISSN: 0031-613X.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
ENTRY DATE: Entered STN: 27 Oct 1997  
Last Updated on STN: 10 Dec 1997

AB The recent characterization of triallate-resistant lines of wild oat (*Avena fatua* L.) deficient in triallate sulfoxidation provides an experimental system to investigate and differentiate the effects of triallate and triallate sulfoxide on wax and lipid biosynthesis. Greenhouse applications of triallate dramatically reduced epicuticular wax deposition in susceptible (S) but not resistant (R) wild oats. Triallate treatment had no effect on in-vivo concentrations of C-12 to C-26 fatty acids and fatty alcohols in R plants, while **elongated fatty acid** fractions (C gt 18) were significantly reduced in S plants. In contrast, treatment with triallate sulfoxide reduced in-vivo concentrations of **elongated fatty acids** equally in R and S, supporting the hypothesis that triallate sulfoxide is more inhibitory than triallate towards fatty acid **elongases**. Although de-novo synthesis of short-chain fatty acids was not affected by triallate or triallate sulfoxide in R or S plants, synthesis of **elongated fatty acid** fractions was dramatically reduced in S plants by triallate. Fatty acid biosynthesis in R and S plants was equally sensitive to triallate sulfoxide. The results support the idea that in-vivo triallate sulfoxidation is necessary for herbicidal activity, and confirm that reduced rates of triallate sulfoxidation confer resistance in R wild oats.

L5 ANSWER 11 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN  
ACCESSION NUMBER: 2005:44368 CAPLUS



DOCUMENT NUMBER: 142:311853  
 TITLE: Biochemical characterization of **elongase** activity in corn (*Zea mays* L.) roots  
 AUTHOR(S): Schreiber, Lukas; Franke, Rochus; Lessire, Rene  
 CORPORATE SOURCE: Department of Ecophysiology, Institute of Cellular and Molecular Botany (IZMB), University of Bonn, Bonn, D-53115, Germany  
 SOURCE: Phytochemistry (Elsevier) (2005), 66(2), 131-138  
 CODEN: PYTCAS; ISSN: 0031-9422  
 PUBLISHER: Elsevier B.V.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Chemical anal. of 4-day-old corn (*Zea mays* L.) root cell walls revealed that the lipophilic biopolymer suberin forms an important constituent of rhizodermal and hypodermal cell walls. Identified aliphatic monomers had chain lengths ranging from C16 to C26 and they belonged to 5 substance classes ( $\omega$ -hydroxycarboxylic acids, 1, $\omega$ -dicarboxylic acids, 2-hydroxycarboxylic acids, carboxylic acids and alcs.) by which suberin is characterized. Biochem. expts. proved the occurrence of **elongase** activities in corn roots. Highest enzymic activities were found in corn root microsomes, and major products synthesized by root **elongases** were **elongated fatty acids** with chain lengths ranging from C20 to C24. Preferred substrates of root **elongases** were acyl-CoAs of the chain length C18 and C20, whereas monounsaturd. acyl-CoAs (C16:1 and C18:1) and acyl-CoAs of lower (C12-C16) and higher chain lengths (C22-C24) were rarely elongated. **Elongase** activities significantly decreased over the length (40 cm) of 10-day-old corn roots going from the young tip to the older base of the root. Thus, results presented here show the presence and activity of **elongases** in roots of plants.

REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 12 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:900207 CAPLUS  
 DOCUMENT NUMBER: 142:89851  
 TITLE: Biosynthesis of very-long-chain polyunsaturated fatty acids in transgenic oilseeds: Constraints on their accumulation

AUTHOR(S): Abbadi, Amine; Domergue, Frederic; Bauer, Joerg; Napier, Johnathan A.; Welti, Ruth; Zaehring, Ulrich; Cirpus, Petra; Heinz, Ernst

CORPORATE SOURCE: Institut fuer Allgemeine Botanik, Universitaet Hamburg, Hamburg, 22609, Germany

SOURCE: Plant Cell (2004), 16(10), 2734-2748  
 CODEN: PLCEEW; ISSN: 1040-4651

PUBLISHER: American Society of Plant Biologists  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The  $\omega$ 6- and  $\omega$ 3-polyunsaturd. C20 fatty acids represent important components of the human diet. A more regular consumption and an accordingly sustainable source of these compds. are highly desirable. In contrast with the very high levels to which industrial fatty acids have to be enriched in plant oils for competitive use as chemical feedstocks, much lower percentages of very-long-chain polyunsaturd. fatty acids (VLCPUFA) in edible plant oils would satisfy nutritional requirements. Seed-specific expression in transgenic tobacco (*Nicotiana tabacum*) and linseed (*Linum usitatissimum*) of cDNAs encoding fatty acyl-desaturases and **elongases**, absent from all agronomically important plants, resulted in the very high accumulation of  $\Delta$ 6-desaturd. C18 fatty acids and up to 5% of C20 polyunsaturd. fatty acids, including arachidonic and eicosapentaenoic acid. Detailed lipid analyses of developing seeds from transgenic plants were interpreted as indicating that, after desaturn. on phosphatidylcholine,  $\Delta$ 6-desaturd. products are immediately channeled

to the triacylglycerols and effectively bypass the acyl-CoA pool. Thus, the lack of available  $\Delta 6$ -desatd. acyl-CoA substrates in the acyl-CoA pool limits the synthesis of **elongated C20 fatty acids** and disrupts the alternating sequence of lipid-linked desaturations and acyl-CoA dependent elongations. As well as the successful production of VLCPUFA in transgenic oilseeds and the identification of constraints on their accumulation, our results indicate alternative strategies to circumvent this bottleneck.

REFERENCE COUNT: 56 THERE ARE 56 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 13 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 2004:557613 CAPLUS

DOCUMENT NUMBER: 141:346651

TITLE: Zebrafish cDNA Encoding Multifunctional Fatty Acid Elongase Involved in Production of Eicosapentaenoic (20:5n-3) and Docosahexaenoic (22:6n-3) Acids

AUTHOR(S): Agaba, Morris; Tocher, Douglas R.; Dickson, Cathryn A.; Dick, James R.; Teale, Alan J.

CORPORATE SOURCE: Institute of Aquaculture, University of Stirling, Stirling, FK9 4LA, UK

SOURCE: Marine Biotechnology (2004), 6(3), 251-261

CODEN: MABIFW; ISSN: 1436-2228

PUBLISHER: Springer-Verlag New York Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Enzymes that increase the chain length of fatty acids are essential for biosynthesis of highly unsatd. fatty acids. The gLELO gene encodes a protein involved in the elongation of polyunsatd. fatty acids in the fungus *Mortierella alpina*. A search of the GenBank database identified several expressed sequence tag sequences, including 1 obtained from zebrafish (*Danio rerio*), with high similarity to gLELO. The full-length transcript ZfELO, encoding a polypeptide of 291 amino acid residues, was isolated from zebrafish liver cDNA. The predicted amino acid sequence of the open reading frame shared high similarity with the **elongases** of *Caenorhabditis elegans* and human. When expressed in *Saccharomyces cerevisiae*, the zebrafish open reading frame conferred the ability to lengthen the chain of a range of C18, C20, and C22 polyunsatd. fatty acids, indicating not only that biosynthesis of 22:6n-3 from 18:3n-3 via a 24-carbon intermediate is feasible, but also that 1 **elongase** enzyme can perform all 3 elongation steps required. The zebrafish enzyme was also able to **elongate** monounsatd. and saturated fatty acids, and thus demonstrates a greater level of promiscuity in terms of substrate use than any **elongase** enzyme described previously.

REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 14 OF 17 CAPLUS COPYRIGHT 2005 ACS on STN

ACCESSION NUMBER: 1997:600786 CAPLUS

DOCUMENT NUMBER: 127:216314

TITLE: Fatty acid and wax biosynthesis in susceptible and triallate-resistant *Avena fatua* L

AUTHOR(S): Kern, Anthony J.; Jackson, Larry L.; Dyer, William E.

CORPORATE SOURCE: Dep. of Plant, Soil and Environ. Sci., Montana State Univ., Bozeman, MT, 59717-0312, USA

SOURCE: Pesticide Science (1997), 51(1), 21-26

CODEN: PSSCBG; ISSN: 0031-613X

PUBLISHER: Wiley

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The recent characterization of triallate-resistant lines of wild oat (*Avena fatua* L.) deficient in triallate sulfoxidn. provides an exptl.

system to investigate and differentiate the effects of triallate and triallate sulfoxide on wax and lipid biosynthesis. Greenhouse applications of triallate dramatically reduced epicuticular wax deposition in susceptible (S) but not resistant (R) wild oats. Triallate treatment had no effect on in-vivo concns. of C12 to C26 fatty acids and fatty alcs. in R plants, while **elongated fatty acid** fractions ( $C > 18$ ) were significantly reduced in S plants. In contrast, treatment with triallate sulfoxide reduced in-vivo concns. of **elongated fatty acids** equally in R and S, supporting the hypothesis that triallate sulfoxide is more inhibitory than triallate towards fatty acid **elongases**. Although de-novo synthesis of short-chain fatty acids was not affected by triallate or triallate sulfoxide in R or S plants, synthesis of **elongated fatty acid** fractions was dramatically reduced in S plants by triallate. Fatty acid biosynthesis in R and S plants was equally sensitive to triallate sulfoxide. Thus, in-vivo triallate sulfoxidn. is necessary for herbicidal activity, and confirm that reduced rates of triallate sulfoxidn. confer resistance in R wild oats.

REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 15 OF 17 LIFESCI COPYRIGHT 2005 CSA on STN

ACCESSION NUMBER: 2005:77355 LIFESCI

TITLE: Zebrafish cDNA Encoding Multifunctional Fatty Acid Elongase Involved in Production of Eicosapentaenoic (20:5n-3) and Docosahexaenoic (22:6n-3) Acids

AUTHOR: Agaba, M.; Tocher, D.R.; Dickson, C.A.; Dick, J.R.; Teale, A.J.

CORPORATE SOURCE: Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, UK; E-mail: d.r.tocher@stir.ac.uk

SOURCE: Marine Biotechnology [Mar. Biotechnol.], (20040600) vol. 6, no. 3, pp. 251-261. ISSN: 1436-2228.

DOCUMENT TYPE: Journal

FILE SEGMENT: Q4

LANGUAGE: English

SUMMARY LANGUAGE: English

AB Enzymes that increase the chain length of fatty acids are essential for biosynthesis of highly unsaturated fatty acids. The gLELO gene encodes a protein involved in the elongation of polyunsaturated fatty acids in the fungus *Mortierella alpina*. A search of the GenBank database identified several expressed sequence tag sequences, including one obtained from zebrafish (*Danio rerio*), with high similarity to gLELO. The full-length transcript ZfELO, encoding a polypeptide of 291 amino acid residues, was isolated from zebrafish liver cDNA. The predicted amino acid sequence of the open reading frame shared high similarity with the **elongases** of *Caenorhabditis elegans* and human. When expressed in *Saccharomyces cerevisiae*, the zebrafish open reading frame conferred the ability to lengthen the chain of a range of C sub(18), C sub(20), and C sub(22) polyunsaturated fatty acids, indicating not only that biosynthesis of 22:6n-3 from 18:3n-3 via a 24-carbon intermediate is feasible, but also that one **elongase** enzyme can perform all three elongation steps required. The zebrafish enzyme was also able to **elongate** monounsaturated and saturated **fatty acids**, and thus demonstrates a greater level of promiscuity in terms of substrate use than any **elongase** enzyme described previously.

L5 ANSWER 16 OF 17 LIFESCI COPYRIGHT 2005 CSA on STN

ACCESSION NUMBER: 2005:42543 LIFESCI

TITLE: Biosynthesis of Very-Long-Chain Polyunsaturated Fatty Acids in Transgenic Oilseeds: Constraints on Their Accumulation

AUTHOR: Abbadi, Amine; Domergue, Frederic; Bauer, Joerg; Napier, Johnathan A.; Welti, Ruth; Zaehring, Ulrich; Cirpus, Petra; Heinz, Ernst

CORPORATE SOURCE: Institut fuer Allgemeine Botanik, Universitaet Hamburg,  
22609 Hamburg, Germany  
SOURCE: Plant Cell, (20041000) vol. 16, no. 10, pp. 2734-2748.  
ISSN: 1040-4651.  
DOCUMENT TYPE: Journal  
FILE SEGMENT: W2  
LANGUAGE: English  
SUMMARY LANGUAGE: English

AB omega 6- and omega 3-polyunsaturated C20 fatty acids represent important components of the human diet. A more regular consumption and an accordingly sustainable source of these compounds are highly desirable. In contrast with the very high levels to which industrial fatty acids have to be enriched in plant oils for competitive use as chemical feedstocks, much lower percentages of very- long-chain polyunsaturated fatty acids (VLCPUFA) in edible plant oils would satisfy nutritional requirements. Seed-specific expression in transgenic tobacco (*Nicotiana tabacum*) and linseed (*Linum usitatissimum*) of cDNAs encoding fatty acyl-desaturases and elongases, absent from all agronomically important plants, resulted in the very high accumulation of delta6-desaturated C18 fatty acids and up to 5% of C20 polyunsaturated fatty acids, including arachidonic and eicosapentaenoic acid. Detailed lipid analyses of developing seeds from transgenic plants were interpreted as indicating that, after desaturation on phosphatidylcholine, delta6-desaturated products are immediately channeled to the triacylglycerols and effectively bypass the acyl-CoA pool. Thus, the lack of available delta6-desaturated acyl-CoA substrates in the acyl-CoA pool limits the synthesis of elongated C20 fatty acids and disrupts the alternating sequence of lipid-linked desaturations and acyl-CoA dependent elongations. As well as the successful production of VLCPUFA in transgenic oilseeds and the identification of constraints on their accumulation, our results indicate alternative strategies to circumvent this bottleneck.

L5 ANSWER 17 OF 17 BIOTECHNO COPYRIGHT 2005 Elsevier Science B.V. on STN  
ACCESSION NUMBER: 1997:27389654 BIOTECHNO  
TITLE: Fatty acid and wax biosynthesis in susceptible and triallate-resistant *Avena fatua* L.  
AUTHOR: Kern A.J.; Jackson L.L.; Dyer W.E.  
CORPORATE SOURCE: A.J. Kern, Dept. of Plant Soil/Envtl. Sciences,  
Montana State University, Bozeman, MT 59717-0312,  
United States.  
SOURCE: Pesticide Science, (1997), 51/1 (21-26), 19  
reference(s)  
CODEN: PSSCBG ISSN: 0031-613X  
DOCUMENT TYPE: Journal; Article  
COUNTRY: United Kingdom  
LANGUAGE: English  
SUMMARY LANGUAGE: English

AN 1997:27389654 BIOTECHNO  
AB The recent characterization of triallate-resistant lines of wild oat (*Avena fatua* L.) deficient in triallate sulfoxidation provides an experimental system to investigate and differentiate the effects of triallate and triallate sulfoxide on wax and lipid biosynthesis. Greenhouse applications of triallate dramatically reduced epicuticular wax deposition in susceptible (s) but not resistant (R) wild oats. Triallate treatment had no effect on in-vivo concentrations of C.sub.1.sub.2 to C.sub.2.sub.6 fatty acids and fatty alcohols in R plants, while elongated fatty acid fractions (C > 18) were significantly reduced in S plants. In contrast, treatment with triallate sulfoxide reduced in-vivo concentrations of elongated fatty acids equally in R and S, supporting the hypothesis that triallate sulfoxide is more inhibitory than triallate towards fatty acid elongases. Although de-novo synthesis of short-chain fatty acids was not affected by triallate or triallate sulfoxide in R or S plants, synthesis of elongated

fatty acid fractions was dramatically reduced in S plants by triallate. Fatty acid biosynthesis in R and S plants was equally sensitive to triallate sulfoxide. The results support the idea that in-vivo triallate sulfoxidation is necessary for herbicidal activity, and confirm that reduced rates of triallate sulfoxidation confer resistance in R wild oats.